www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023

ORGANIC-MINERAL FERTILIZERS BASED ON ACIDIFIED SEWAGE SLUDGE AND GULIOB PHOSPHORATE

A.A.Mamataliyev – Doctor of Technical Sciences, Institute of General and Inorganic Chemistry, Tashkent
S.E.Qorayev – teacher, Denov Institute of Entrepreneurship and Pedagogy, Denov
E.F.Murodova - student, Denov Institute of Entrepreneurship and Pedagogy, Denov
Z.B.Eshquvatova - student, Denov Institute of Entrepreneurship and Pedagogy, Denov

[1] Denov Institute of Entrepreneurship and Pedagogy, 190507 Denau Sharof Rashidov 360,Republic of Uzbekistan
[2]Institute of General and Inorganic Chemistry, 77a Mirzo Ulugbek, 100170 Tashkent,Uzbekistan

Abstract.The article shows that it is possible to obtain organomineral fertilizers by adding Guliob phosphorite (GPh) by weight ratio SS: GPh = 100 : 2 to 100 : 25 to the municipal sewage sludge (SS) obtained on the basis of acidification in the presence of sulfuric acid in the range of pH 2-4. The optimal ratios of starting materials for the production of organomineral fertilizers and the amount of $P_2O_{5tot.}$, $P_2O_{5assi.}$, $CaO_{tot.}$, CaO_{water}, humic and fulvic acids, water-soluble organic substances in the obtained fertilizers were determined.
Keywords:urban sewage sludge water (SSW), Guliob phosphorite (GPh), sulfuric acid, phosphorus, calcium, humic acid, organomineral fertilizers.

Introduction. The rapid growth of the world's population from year to year and the sharp decline in arable land is exacerbating the global food problem. In this regard, one of the effective solutions to the food problem is the rational use of agrochemically effective mineral fertilizers to increase soil fertility, high and quality yields of agricultural crops. Therefore, it is important to obtain complex fertilizers that contain various nutrients necessary for plant development. This is due to the fact that in addition to macro-and micronutrients necessary for plants, complex fertilizers also contain humic substances that improve soil structure and fertility [1-5].

Among safe substances promising from the point of view of agro chemistry, compounds of natural origin, in particular, humic substances deserve special attention. Their formation (humification) is the second largest process of transformation of organic matter in the environment after photosynthesis, which involves a huge amount of carbon per year.

Humic substances are complex mixtures of biodegradation-resistant high-molecular dark-colored organic compounds of natural origin, formed during the decomposition of plant and animal residues under the action of microorganisms and abiotic environmental factors. Humic substances are a macro component of the organic matter of soil and water ecosystems, as well as solid fossil fuels.

Traditional sources of humic substances include peat, lignin, black cattle and poultry manure, as well as naturally oxidized coals and others.

It is known from the literature that peat bog plants, the main part of which consists of humus substances that increase soil structure and productivity, were formed during natural decay and natural decay in the presence of excess moisture. However, due to the lack of forests and wetlands in Uzbekistan, peat is almost non-existent [6].

However, due to the lack of forests and wetlands in Uzbekistan, peat is almost non-existent. Therefore, a wide range of technologies for the production of organic mineral fertilizers based on livestock, poultry waste and Kyzylkum phosphorites, oxidized Angren brown coal,

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023

ammophos, suprefos, urea and phosphogypsum waste from JSC "Ammophos-Maxam" has been developed [7-12].

Other sources rich in humic substances can also include municipal sewage sludge (SS). The processes of obtaining organomineral fertilizers on the basis of SS were studied and it was found that the application of complex fertilizers to soils and crops not only increased their productivity, but also significantly reduced the cost of organomineral fertilizers obtained compared to other complex fertilizers [13-16].

Raw materials for obtaining organic and organomineral fertilizers include: bedding manure, bedding-free manure, bird droppings, peat, brown coal, green manure, straw, sapropel, household and industrial waste, as well as SSW [17-19].

In the patent [20], heavy metal ions are first removed from sewage sludge by treating urban sewage sludge (SSW) with a solution of nitric acid with a concentration of 1.0-1.25 mol/dm3 at a temperature of 50-70°C for 10-20 minutes. The process is carried out at a ratio of solid and liquid phases equal to 1: 5. Under these conditions, the residual content of heavy metals corresponds to the standards for this type of fertilizer. After filtration, the residual acidity is neutralized with alkaline agents such as potassium hydroxide, ammonium hydroxide, calcium carbonate to obtain an organomineral fertilizer. The metals isolated from the sediment are precipitated in the solution and processed into sludge, from which they are then regenerated.

In next patent [21], 1000 l of a mixture is prepared from 200 l of SS water in a container which the mixture is circulated. The temperature of the prepared mixture is 40°C. Then, an alkaline reagent - 12.5 kg of NaOH is added to the container by the prepared mixture, and the mixture is alkalinity extracted with ultrasonic treatment for 60 minutes in an ultrasonic reactor. Also, the mixture is settled for a day to separate the ultrafine suspension of fibrous organic material, the liquid phase is drained and the insoluble part of the sediment is unloaded. As a result, 900 liters of liquid organomineral fertilizer are obtained and about 100 liters of the sediment of the latter are used as fertilizer for growing industrial and forest crops.

Materials and methods. However, in scientific studies based on SSW, the microorganisms that cause pathogenic diseases in it are not completely deactivate. Therefore, in our opinion, it was considered expedient to obtain complex organomineral fertilizers based on inorganic additives containing SSW, which are useful elements for plants. Termez city SSW (wt.%) Moisture - 65.43; ash - 9.74; organic matter - 24.62; humic acids - 4.35; fulvic acids - 7.87; water-soluble organic substances - 4.23; P_2O_5 - 1.43; N 1.29; K_2O - 0.53; CaO - 4.31, sulfuric acid produced in Uzbekistan and phosphorite from the Guliob deposit in Sariosiya district, Surkhandarya region were selected.

Table 1 shows the chemical and dispersion compositions of Guliob phosphorite.

Table 1

Chemical and disperse composition of Guliob phosphorite (GPh)

Content of components, wei. %		Class size, mm	Fraction yield, wt. %
P_2O_5	7,88	- 2 + 0,5	16,1
CaO	20,64	- 0,5 + 0,315	52,3
Al_2O_3	0,62	- 0,315 + 0,16	13,4
Fe_2O_3	0,45	- 0,16 + 0,1	3,5

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023

MgO	1,02	- 0,1 + 0,063	9,2
F	1,36	- 0,063 + 0,05	3,9
CO_2	8,03	- 0,05	1,6
$P_2O_{5assi.}$ for trilon B $P_2O_{5tot.}$, %	21,32	Exhaust mass	100

From Table 1 shows that GPh is characterized by a low content of phosphorus (7.88% P_2O_5), a high content of carbonates (20.64% CO_2) and a high value of the calcium module (CaO : P_2O_5 = 5.2). Chemical analysis of the SSW of the city of Termiz, GPh and products of their processing was carried out by the methods below. Humidity was determined according to GOST 26712-85, ash content according to GOST 26714-85 and organic matter according to GOST 27980-80.

The amount of the water-soluble fraction of organic substances extracted from the products with water was determined by filtration and evaporation on a water bath, drying the solid residue to a constant weight, followed by burning it to determine the ash content and subtract it. Humic acids were isolated by treating the products with a 0.1 n alkali solution, followed by acidification of the solution with mineral acid [22].

The solid phase after the separation of alkali-soluble organic substances from it contains residual organic matter. It was thoroughly washed with distilled water, dried to a constant weight, and the content of organic substances was determined. The difference between the amounts of alkali-soluble organic substances and humic acids gives us the content of fulvic acids. All forms of P_2O_5 were determined by the gravimetric method by precipitating the phosphate ion with a magnesia mixture in the form of magnesium ammonium phosphate, followed by calcining the precipitate at 1000-1050°C according to GOST 20851.2-75. Digestible forms of P_2O_5 were determined by solubility both in 2% citric acid and in 0.2 M Trilon B solution. CaO content was determined complexometrically : by titration with 0.05 N Trilon B solution in the presence of fluorexone indicator. 92% H_2SO_4 was used to acidify SSW.

The experiments were carried out as follows: first the measured sulfuric acid was poured into a glass reactor with a volume of 5 l, and then a certain amount of measured SSW was gradually added to it. Sulfuric acid was acidified by adding SSW: H_2SO_4 = 100:10 to 100:50 by weight relative to the organic component in SSW. In the last stage of the experiment, complex type organomineral fertilizers were obtained by adding to the acidified SSW slurry: GPh = 100 : 10 to 100 : 50 by weight. The experiments were performed at 25°C temperature and for 60 min.

Results and discussion. The results obtained are shown in Table 2 and Figures 1-4.

Table 1

Composition of complex fertilizers based on SSW sulfuric acid and GPh

Weight ratio SS: GPh	**$P_2O_{5tot.}$**	**$P_2O_{5assi.}$**	**$P_2O_{5wat.}$**	**$CaO_{tot.}$**	**$CaO_{assi.}$**	**CaO_{wat}**	**Organic**	**Humic**
Weight ratio SS organic : H_2SO_4 = 100 : 10								
100 : 10	3,94	1,95	0,21	13,88	8,26	1,72	44,53	29,75

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023

100 : 20	4,47	2,07	0,20	18,40	10,50	2,55	37,13	24,81
100 : 30	4,84	2,10	0,19	21,60	12,07	3,14	31,77	21,23
100 : 40	5,12	2,11	0,16	23,97	13,10	3,72	27,75	18,54
100 : 50	5,33	2,12	0,13	25,78	13,59	4,29	24,60	16,44
Weight ratio SSW organic : H_2SO_4 = 100 : 20								
100 : 10	3,71	2,21	0,23	13,08	8,57	3,24	41,97	28,04
100 : 20	4,28	2,38	0,24	17,60	10,92	4,78	35,51	23,73
100 : 30	4,68	2,47	0,22	20,86	12,62	5,93	30,69	20,50
100 : 40	4,98	2,49	0,19	23,32	13,84	6,92	26,99	18,03
100 : 50	5,19	2,50	0,15	25,14	14,20	7,96	23,99	16,03
Weight ratio SSW organic : H_2SO_4 = 100 : 30								
100 : 10	3,51	2,50	0,27	12,35	9,01	4,35	39,63	26,48
100 : 20	4,09	2,73	0,26	16,82	11,94	6,34	33,93	22,67
100 : 30	4,52	2,86	0,25	20,13	14,12	7,90	29,62	19,79
100 : 40	4,83	2,89	0,22	22,64	15,61	9,11	26,21	17,51
100 : 50	5,08	2,92	0,18	24,60	16,35	10,46	23,47	15,68
Weight ratio SSW organic : H_2SO_4 = 100 : 40								
100 : 10	3,32	2,70	0,31	11,68	10,00	5,33	37,49	25,05
100 : 20	3,91	3,13	0,30	16,08	12,88	7,65	32,44	21,67
100 : 30	4,35	3,42	0,29	19,39	14,76	9,34	28,52	19,06
100 : 40	4,68	3,61	0,26	21,94	15,56	10,88	25,39	16,97
100 : 50	4,95	3,70	0,21	23,96	16,33	12,62	22,86	15,28
Weight ratio SSW organic : H_2SO_4 = 100 : 50								
100 : 10	3,15	2,82	0,36	11,08	10,84	6,03	35,56	23,76
100 : 20	3,73	3,23	0,35	15,36	14,58	8,73	30,99	20,70
100 : 30	4,18	3,50	0,34	18,64	17,02	10,86	27,41	18,32
100 : 40	4,53	3,70	0,30	21,22	18,06	12,70	24,56	16,41
100 : 50	4,80	3,78	0,24	23,21	18,99	14,34	22,15	14,80

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023




Figure 1. $P_2O_{5assi.}$ dependence of the transition to the weight ratios of SSW organic: H_2SO_4 and SSW: GPh.




Figure 2. $P_2O_{5wat.}$ dependence of the transition to the weight ratios of SSW organic: H_2SO_4 and SSW: GPh.

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023




Figure 3. $CaO_{assi.}$ dependence of the transition to the weight ratios of SSW organic: H_2SO_4 and SSW: GPh.




Figure 4. $CaO_{wat.}$ dependence of the transition to the weight ratios of SSW organic: H_2SO_4 and SSW: GPh.

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023

It can be seen from them that $P_2O_{5tot.}$ as the weight ratio of GPh added to the acidified SSW porridge increases. $P_2O_{5assi.}$ $CaO_{tot.}$ $CaO_{assi.}$ $CaO_{wat.}$ while the soluble forms were found to be increased, the organic content of SSW as well as the humic content was found to be reduced. For example, SSW organic: H_2SO_4 = $P_2O_{5tot.}$ in products with the addition of Guliob phosphorite in a ratio of 100 : 10 to 100 : 50 to the fermented SSW porridge in the ratio of 100 : 10 by weight. $P_2O_{5assi.}$ $CaO_{tot.}$ $CaO_{assi.}$ $CaO_{wat.}$ forms from 3.94 to 5.33%, respectively; 1.95 and 2.12% respectively; 13.88 and 25, 78% respectively; 8.26 and 13.59% respectively; Organic content and humus content decreased from 44.53 to 24.60% and from 29.75 to 16.44%, respectively, while increasing from 1.72 to 4.29%.

The best performance in this study was found in products with the addition of Guliob phosphorite in a ratio of 100 : 10 to 100 : 50 to the acidified SSW slurry in the ratio of SSW organic : H_2SO_4 = 100 : 50 by weight. Here $P_2O_{5tot.}$ $P_2O_{5assi.}$ $CaO_{tot.}$ $CaO_{assi.}$ $CaO_{wat.}$ forms from 3.15 to 4.80%, respectively; 2.82 and 3.78% respectively; 11.08 and 23.21% respectively; 10.84 and 18.99% respectively; Organic content and humus content decreased from 35.56 to 22.15% and from 23.76 to 14.80%, respectively, while increasing from 6.03 to 14.34%. $P_2O_{5tot.}$ in the composition of phosphorus fertilizers used in agricultural crops. No matter how high the shape, the $P_2O_{5assi.}$ forms requires no less than 50%. In almost all organomineral fertilizers obtained by us, it was found that the $P_2O_{5assi.}$ form is not less than 50%. In Figures 1-4, the plant-replicating $P_2O_{5tot.}$ $P_2O_{5wat.}$ and $CaO_{wat.}$ $CaO_{wat.}$ the increase in soluble forms is evident.

Conclusions. It can be concluded from the results of scientific research that organomineral fertilizers can be obtained first by acidifying SSW with sulfuric acid and then by adding Guliob phosphorite to the acidified SSW slurry. It was found that the elements of phosphorus and calcium in plants containing Guliob phosphorite are absorbed by inorganic (sulfuric acid) and organic (humic and fulvic acids) acids.

References:

[1] Sh.Namazov, U. Temirov, N. Usanbayev Research of the process of obtaining organo-mineral fertilizer based on nitrogen acid decomposition of non-conditional phosphorites of central kyzylkumes and poultry cultivation waste. International Journal of Innovative Technology and Exploring Engineering (IJITEE), V.8. No. 12. October 2019 PP. 2260-2265.

[2] Rasulov O, Mamataliyev A, Rasulova D, Temirov U, Namazov Sh 2021 Physico-chemical properties lime-ammonium nitrate based on chalk, nitrate and ammonium sulphate E3S Web of Conferences 264 04012.

[3] Temirov U, Doniyarov N, Jurakulov B, Usanbaev N, Tagayev I, Mamataliyev A 2021 Obtaining complex fertilizers based on lowgrade phosphorites E3S Web of Conferences 264 04009.

[4] Sh. Namazov, U. Temirov, N. Usanbayev Intensive technology for processing bird litter in organomineral fertilizers. Izv. Vyssh. Uchebn. Zaved. Khim. Khim. Tekhnol. [Russ. J. Chem. & Chem. Tech.]. 2020. V. 63. N 12. P. 85-94.

[5] Zhumanova M.O., Sh. Namazov, B. Beglov, O. Myachina, M. Tashquziev Influence of organic and organic-mineral fertilizers on the fertility of soils. Sofia. Journal of Chemical Technology and Metallurgy (2015) V.50. N.3.P. 282-287.

[6] Anders Ekbom, Yonas Alem and Thomas Sterner. Integrating soil science into agricultural production frontiers. Environment and Development Economics. V. 18. No. 3 (June 2013), pp. 291-308.

[7] Derzhavin L.M., Polyakov A.N., Florinsky M.A. Humus content in arable soils of the USSR. Chemicalization of agriculture. 1988. No. 6. P. 7-8.

[8] Ivanov V.M. Humus: basics of fertility. AgroONE (Ukraine). 2018. No. 24. P.12-13.

www.wordlyknowledge.uz Volume: 3, Issue: 1, 2023

[9] N.G. Kovalev Modern problems of production and use of organic fertilizers. Bulletin of the All-Russian Research Institute of Livestock Mechanization. 2013 No. 2(10). P.82-92.
[10] Titova I.N. Humates and soil. Moscow: ILKO, 2006. - P.9-12.
[11] Derzhavin L.M., Sedova E.V. On the issue of humus reproduction. Agrochemistry. 1988. No. 9. P. 117-127.
[12] Rasulov O, Mamataliyev A, Rasulova D, Temirov U, Namazov Sh 2021 Physico-chemical properties lime-ammonium nitrate based on chalk, nitrate and ammonium sulphate E3S Web of Conferences 264 04012. Scopus (3) IF – 0.203.