AGROTECHNOLOGY OF MAIZE CULTIVATION AND A BRIEF CLASSIFICATION OF VIRAL DISEASES

G'afforova Gavhar

Introduction

Rural property in ensuring food safety in the country getting a high, high-quality harvest from their crops, fighting the diseases of the ulninig,conducting preventive measures is one of the most pressing problems today one. In meeting our population's demand for meat and dairy products the corn plant is incredibly large in food importance it is distinguished by its ownership.

Corn- " Hori Zea mays L. - an annual belonging to the Virgo family the plant.is. The homeland of the choir"Corn" is central and South America the region is. The growing period is 90-150 depending on the varieties and climatic conditions of the soil consists of a day. It sprouts in 10-12 days when the soil temperature reaches 10°C. 20-25°C develops normally.Corn's personality is widely seen in folk culture used. In many countries on Earth, 20 per cent of the maize grain used as food. Flour, cereals, preserves are made from it. In industry starch, ethyl alcohol, dextrin, glucose, ascorbic acid and glutamine acids are obtained.

The columnar part of the flower is used in the medicine. Corn "Chori so" used from stem and leaf, linolium, viscose materials are taken. In our country, mackerel is a crop from which silage is prepared as it stands in the first place. Cereals contain 9-12 percent protein, fat" 4-8 percent,carbohydrates 65-70 percent also contain cultured salts and vitamins. The STEM is the feed is used as a hash. The macajoo crop is the most cultivated in the world one of the cereal plants is planted on more than 135 million hectares in the globe.In our country, a large number of hundred centners are harvested in advance.High yields are obtained by planting in all districts of Uzbekistan.

Importance in crop rotation. One pitch to the McCaughey"choir being resistant to repeated plantings, the corn choir in Uzbekistan is mainly sugarcane is cultivated on cultivated land. It can be grown in alfalfa, germ, potatoes, pulses, autumn cereals placement after crops gives a good result.Corn choir after lavlagí when placed phosphates are transferred, nutritional conditions worsen.The placement of the "corn" choir after grain crops also increases productivity, grain improves quality.

The yield is 3-4 ts after the macajoo with respect to monoculture increased. The placement of the corn " choir in crop rotation is a unit of food from one hectare increases the efficiency of irrigated land by increasing the output.Corn " choir autumn wheat, autumn barley, potato is a good native crop for melons is. In a field to repel him for 4-5 years, in non-karakuya areas 10-15 years can be transferred. This crop was taken by the farm in crop rotation, to the farm planting in nearby areas, increasing organic growth, harvesting, it is possible to reduce storage, transportation work, costs gives.

Uneven fields are leveled before working the soil. Soil the main method of operation and the depth of the property of the parent crop, soil culture to the thickness of the layer, the degree of pressing the field, the type of soil, last year it is determined depending on the depth of expulsion. Fields of interest if the driving layer is thick, 25-28 CM, some years fall to a depth of 40-45 CM plowed. The soil is 40-45 CM with two-slit plugs with a pinch of a year if recorded in depth, the second year is 25-28 cm deep, and the third year is another 40-45 SM is driven to the depth. So the eviction system is applied to the remnants of the anchor,weed seeds, pests buds, sources of disease for two years during rotting, neutralizing, soil fertility increases goes. Autumn plow in the northern region of our republic from October 20-30 to October 30 it will be completed until November, 15th in the southern regions. Newly acquired the soils are 20-22 cm deep in the first year and 2 -3 CM annually in the following

years deepened and expelled. Mechanical composition heavy, dense plastered layer 40-50 CM soils located in depth, special at a depth of 40-50 CM before driving it is loosened with tools, then driven to a depth of 25-30 CM. Strong plowing the infested fields with two-wound plugs to a depth of 35-40 CM, alien reduces the yield of grain compared to driving at a depth of 27 cm ts / Ga increases. If the soil is dry before plowing, the field before plowing sucked. Bedapoyas 5-8 with lushchilnik or otvalless plugs before driving the heads of alfalfa beds driven to a depth of CM are cut off. 30 after a week-Driven to a depth of 40 cm. Driving depth 20-22 CM in the second year, third the year is 30-40 cm. Fields from grain crops with lushchilniks 6-8 CM it is loosened, then driven to a depth of 25-27 CM. Soil maturation in early spring baronage with is conducted.

Conclusion

Usually in the first place almost on the lower parts of the youngest leaves colorless will appear in the form of more or less rounded spots. Then the lesion it covers the entire base of the leaf, grows on other leaves and has veins it takes the form of spaced chlorotic lines. The length of the lines is one varies from a few millimeters to a few centimeters.

REFERENCES:

1. Mutalov K.A., Ramazonov B.R., Boqiyev D.T., Ishmo"minov B.B. Laboratory on the subject of biological foundations of"Soil Science and rural education" and practical exercises " training. Tashkent:” reliable partner", 2021. – 168PP.

2. Atabayeva H. Qadirkhujayev O. Academic, T.Generation of the new century. 2006 (in the Latin alphabet).

3. Yormatova D.Practical from field crop biology and cultivation technology fame.

4. .Azimboev S.A. Fundamentals of Agriculture, soil science and agrochemistry.

5. Mutalov K., Pamazonov B, Zakirov D, Gulmatova M. Polukustarnichkovaya rastitelnost partekovaya formasiya convolvuleta hamadae. academic research in educational sciences. 2020(2):205-12.