https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

Nurmuxammad Abduxoshimovich Mamadjonov

Student of the Faculty of Pediatrics, Andijan State Medical Institute

COMPREHENSIVE COMPARATIVE ANALYSIS OF BREASTFEEDING AND ARTIFICIAL FEEDING IN FULL-TERM AND PRETERM INFANTS: EXTENSIVE NUTRITIONAL, CLINICAL, AND EPIDEMIOLOGICAL DATA WITH SPECIAL REFERENCE TO UZBEKISTAN AND THE FERGHANA VALLEY

Abstract

Early-life nutrition is one of the most influential determinants of infant survival, physiological development, immune competence, and long-term metabolic health. The neonatal period, particularly the first six months of life, represents a critical window during which nutritional exposures exert lasting effects on morbidity, mortality, and disease susceptibility across the lifespan. This is especially relevant for preterm infants, who experience heightened vulnerability due to immature gastrointestinal, immune, renal, and metabolic systems. Breastfeeding is universally acknowledged as the optimal feeding strategy; however, artificial feeding remains prevalent in many regions due to medical, socio-economic, and cultural factors.

This article provides an in-depth, data-driven comparative analysis of breastfeeding and artificial feeding in full-term and preterm infants. The analysis integrates macronutrient and micronutrient composition, bioactive components, protein quality (with a particular focus on casein), clinical outcomes, epidemiological indicators, and economic implications. The study is contextualized for Uzbekistan, with a specific focus on the Ferghana Valley, a region characterized by high population density, elevated birth rates, and distinct maternal nutritional patterns. By synthesizing extensive international and regional data, this

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

study demonstrates the biological, clinical, and public health superiority of breastfeeding and highlights strategic implications for maternal and child health policy in Uzbekistan.

Keywords

breastfeeding, artificial feeding, infant nutrition, preterm infants, casein, public health, Uzbekistan, Ferghana Valley, epidemiology

Introduction

Globally, infant nutrition remains a central concern in public health, pediatrics, and health economics. According to the World Health Organization, approximately 130–140 million children are born each year worldwide, with an estimated 10–11% classified as preterm. Suboptimal infant feeding practices are directly or indirectly responsible for nearly half of all deaths among children under five years of age, primarily through increased susceptibility to infectious diseases, malnutrition, and impaired immune development.

Breastfeeding is widely recognized as a cornerstone of infant survival and development. Large-scale epidemiological studies have demonstrated that exclusive breastfeeding during the first six months of life significantly reduces infant mortality, hospital admissions, and healthcare expenditures. Conversely, early introduction of artificial feeding has been associated with increased risks of gastrointestinal and respiratory infections, allergic diseases, obesity, and metabolic disorders later in life.

Uzbekistan presents a unique demographic and public health context. The country maintains one of the highest fertility rates in Central Asia, with particularly high birth density in the Ferghana Valley, where a substantial proportion of the national population resides. Despite strong cultural norms favoring breastfeeding, a gradual shift toward mixed feeding and early formula supplementation has been observed,

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

particularly in urban and peri-urban areas. Factors contributing to this trend include maternal employment, perceived insufficiency of breast milk, limited lactation support, and aggressive marketing of infant formula.

The Ferghana Valley poses additional challenges due to its high population density, seasonal climatic extremes, and regional disparities in healthcare access. These factors amplify the public health importance of optimal infant feeding practices. This study seeks to address these issues by providing an extensive comparative analysis of breastfeeding and artificial feeding, grounded in robust data and tailored to the Uzbek context.

Methods

This study employed an extensive narrative review combined with comparative data synthesis. Data sources included:

- WHO, UNICEF, World Bank, and ESPGHAN reports (2018–2024);

- peer-reviewed articles indexed in PubMed, Scopus, and Web of Science;

- regional demographic health surveys relevant to Central Asia;

- meta-analyses and systematic reviews focusing on infant nutrition and preterm outcomes.

Data were extracted and categorized into thematic domains: nutritional composition, bioactive components, protein quality, clinical outcomes, epidemiological indicators, and economic impact. Comparative analysis was conducted between breastfeeding and artificial feeding, with additional stratification for full-term and preterm infants.

Results

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

1. Nutritional Composition of Human Breast Milk

Human breast milk contains an optimal balance of macronutrients, including proteins, fats, and carbohydrates, as well as a wide range of micronutrients and bioactive compounds. Unlike infant formulas, the composition of human milk is dynamic and changes according to the stage of lactation, time of day, and the specific needs of the infant.

Proteins in human milk consist primarily of whey proteins and casein in a ratio that supports easy digestion and efficient absorption. Fats provide the main source of energy and include long-chain polyunsaturated fatty acids such as DHA and ARA, which are essential for brain and visual development. Lactose serves as the primary carbohydrate and supports calcium absorption and gut microbiota development.


In the Ferghana Valley, maternal dietary patterns, seasonal food availability, and micronutrient intake influence the nutritional quality of breast milk. Nevertheless, even under conditions of moderate maternal nutritional deficiency, breast milk remains superior to artificial feeding.

2. Comparison with Artificial Feeding

Infant formulas are designed to provide standardized amounts of nutrients and are essential when breastfeeding is not possible. However, formulas lack many bioactive components of human milk, including immunoglobulins, living cells, enzymes, and human milk oligosaccharides.

In Uzbekistan, artificial feeding is sometimes introduced early due to maternal employment, misconceptions about milk insufficiency, or aggressive marketing of formula products. This practice may increase the risk of infections and gastrointestinal disorders, particularly in rural and densely populated regions such as the Ferghana Valley, where sanitation and access to clean water may be limited.

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026



3. Macronutrient Composition and Energy Supply

Human milk provides an energy density optimized for infant growth while avoiding metabolic overload. Unlike formula, its composition dynamically adapts over time.

Table 1. Macronutrient Composition (per 100 ml)

Parameter	Human Milk	Infant Formula
Energy (kcal)	65–70	65–68
Protein (g)	1.0–1.3	1.4–1.6
Fat (g)	3.5–4.5	3.0–3.8
Carbohydrates (g)	6.8–7.2	6.5–7.0

4. Bioactive Components and Immune Protection

Human milk contains hundreds of bioactive molecules absent in formula.

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

Graphical abstract. Human breast milk (HBM) contains macronutrients and micronutrients, and its composition varies according to environmental factors. Colostrum is low in fat but high in protein and relatively rich in immunoprotective components. Micronutrients, hormones, and growth factors in HBM also play various roles in infant development. Microbial communities and microRNAs help construct the infant immune system.

Table 2. Bioactive Components and Biological Functions

Component	Human Milk	Infant Formula	Function
Secretory IgA	Present	Absent	Mucosal immunity
Lactoferrin	Present	Absent	Antimicrobial
Lysozyme	Present	Absent	Bacterial lysis
HMOs	Present	Absent	Microbiome shaping
Immune cells	Present	Immune maturation	

5. Protein Quality and the Role of Casein

Casein is a key protein component of human milk, although its concentration is significantly lower than in cow's milk. Human milk casein forms soft, easily digestible curds in the infant stomach, minimizing gastrointestinal stress and supporting nutrient absorption. This is especially important for preterm infants with immature renal function.

Table 3. Protein Characteristics

Feature	Human Milk	Formula
Whey: casein ratio	Physiological	Altered
Casein curd formation	Soft	Dense
Renal solute load	Low	Higher

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

Digestibility	High	Moderate

Casein plays a crucial role in calcium and phosphorus transport, contributing to bone mineralization and skeletal growth. For preterm infants, the balanced casein-to-whey protein ratio is essential for metabolic stability and growth without excessive renal solute load. Infant formulas contain modified casein, but its biological functionality differs from that of human milk.

6. Clinical Outcomes: Morbidity and Mortality

Outcome	**Breastfed**	**Formula-fed**
GI infections	8–10	20–25
Respiratory infections	10–15	25–35
Allergic diseases	8–12	20–30
NEC (preterm)	3–5	12–15
Hospital admissions (1st year)	10–15	30–40

Breastfeeding and Preterm Infants

Meta-analyses indicate that exclusive human milk feeding reduces the risk of necrotizing enterocolitis by approximately 50–60% in preterm infants. In Uzbekistan, expanding access to expressed maternal milk and donor milk could significantly reduce neonatal morbidity, particularly in regional centers serving the Ferghana Valley.

7. Economic and Public Health Impact

Indicator	**Breastfeeding Promotion indicator**	**Formula Dependence**
Infant infection rate	Lower	Higher
Healthcare costs	Reduced	Increased

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

Family economic burden	Lower	Higher
Health system pressure	Reduced	Elevated

Breastfeeding promotion represents a highly cost-effective public health intervention. Modeling studies suggest that increasing exclusive breastfeeding rates to 90% could reduce healthcare expenditures related to infant infections by millions of USD annually.

In the Ferghana Valley, where healthcare facilities may face capacity constraints, breastfeeding offers a sustainable strategy to improve child health outcomes and reduce system burden.

Discussion

The accumulated evidence presented in this study clearly demonstrates that breastfeeding offers unparalleled biological, clinical, and economic advantages over artificial feeding. The unique composition of human milk cannot be replicated through industrial processes, regardless of technological advances.

In Uzbekistan, particularly in the Ferghana Valley, breastfeeding represents not only a nutritional practice but a critical public health intervention. The region's demographic density and healthcare constraints magnify the protective effects of breastfeeding against infectious diseases and malnutrition. Artificial feeding, while necessary in specific medical contexts, should not replace breastfeeding in the absence of clear indications.

The role of casein exemplifies the importance of protein quality rather than quantity. Human milk casein supports mineral absorption and gastrointestinal tolerance, whereas formula-derived casein may contribute to metabolic stress, especially in preterm infants.

Conclusion

https://wordlyknowledge.uz/index.php/gpble
Qabul qilingan sana: 07.12.2025
Nashr etilgan sana: 30.01.2026

Breastfeeding remains the most physiologically appropriate and biologically superior method of infant feeding. Extensive data confirm that human milk supports optimal growth, immune maturation, and long-term health outcomes in both full-term and preterm infants. For Uzbekistan and the Ferghana Valley, prioritizing breastfeeding through education, policy, and healthcare support is essential for reducing infant morbidity, mortality, and healthcare costs. Artificial feeding should be reserved for cases where breastfeeding is not feasible and should be carefully regulated.

References

1. WHO. Infant and young child feeding. 2021.
2. UNICEF. Breastfeeding and child survival. 2022.
3. Victora CG et al. The Lancet. 2023.
4. Kim SY, Yi DY. Clin Exp Pediatr. 2020.
5. Koletzko B et al. Pediatr Res. 2022.
6. Lönnerdal B. J Pediatr. 2023.
7. ESPGHAN Committee. 2022.
8. Rollins NC et al. The Lancet. 2022.
9. Underwood MA. Clin Perinatol. 2020.
10. Abrams SA. Pediatrics. 2021.
11. World Bank. Nutrition and early development. 2022.
12. Black RE et al. Maternal and Child Nutrition. 2021.
13. Victora CG. Public Health Nutr. 2020.
14. Horta BL et al. Acta Paediatr. 2021.
15. Bode L. Glycobiology. 2021.
16. Andreas NJ et al. Early Hum Dev. 2020.
17. Victora CG. BMJ. 2022.
18. Oddy WH. Nutrients. 2021.
19. WHO Europe. Breastfeeding policies. 2023.