IQRO JURNALI / 2023
GUVOHNOMA № 060680
E-ISSN : 2181-4341
https://wordlyknowledge.uz
IMPACT FACTOR(SJIF) : 3,934

IMPACT FACTOR (RESEARCH BIB) – 7,245
SPECIAL ISSUE, VOLUME 5, ISSUE 2, 2023

PNEUMONIA IN CHILDREN UP TO SCHOOL AGE

Fozilova Nodira Mirkamol qizi,
Gulomova Farangiz Sarvar qizi,
Tosharova Munisa Abdusattor qizi,
Sharapova Kumush Orif qizi.
3rd year student of the Faculty of Pediatrics of Samarkand Medical University

Abstract

Pneumonia remains one of the leading causes of mortality and morbidity in children under five globally. According to WHO estimates, there are over 150 million cases of childhood pneumonia annually, resulting in over 800,000 deaths. Young children, particularly those under two, and those from developing countries face the highest risk. Pneumonia in this vulnerable population warrants focused attention given significant knowledge gaps that hamper effective prevention and clinical management. This article presents a comprehensive review of the current evidence base on childhood pneumonia up to school age. It systematically examines various epidemiological, etiological, clinical, diagnostic and management aspects of the disease. Key findings from the review point to the diversity of risk factors involved, including malnutrition, indoor pollution, lack of exclusive breastfeeding and suboptimal vaccination levels. Prevailing etiological agents like Streptococcus pneumoniae, respiratory syncytial virus and Haemophilus influenzae are discussed. Common clinical features and treatment guidelines endorsed by organizations such as WHO are outlined. Challenges with diagnostic certainty are highlighted, underscoring opportunities for low-cost point-of-care testing. The implications of antibiotic resistance are also explored. Overall, the article aims to consolidate available knowledge in order to inform targeted strategic interventions against pneumonia, one of the leading child health threats globally.

Keywords

Childhood pneumonia, risk factors, epidemiology, etiology, clinical presentation, diagnosis, treatment guidelines, case management, prevention strategies, developing countries, public health.

Pneumonia constitutes a significant disease burden impacting child health worldwide. According to WHO, it accounts for over 15% of all deaths in children under five globally (Rudan et al., 2008). Pneumonia mainly affects young children and considerable geographic inequities exist, with over 95% deaths concentrated in developing countries (Liu et al., 2015). Despite advances in prevention and treatment, pneumonia continues to undermine progress towards child mortality reduction goals espoused by the Sustainable Development Agenda (Alecrim et al., 2019). Given its widespread prevalence and associated morbidity, pneumonia in early childhood necessitates sustained multidisciplinary investigation and policy attention.

This review endeavor seeks to provide a comprehensive synthesis of current knowledge regarding pneumonia in children below school age, up to five years. It systematically examines prevailing epidemiological trends, characterized risk factors, the scope of prevalent etiological agents and major

clinical presentations. Diagnostic approaches and recommended treatment guidelines endorsed by agencies such as WHO are outlined. Gaps are identified to stimulate future research directions. The implications of emerging issues like antibiotic resistance are explored. By consolidating evidence from global sources, the aim is to inform strategic planning and interventions against childhood pneumonia, a pressing public health challenge.

The methodology employed for the review is described in the following section. Primary sources such as peer-reviewed literature and technical reports published over the past two decades have been analyzed. Regional insights are provided to underscore context-specific nuances. By presenting a multifaceted analysis integrating epidemiological, clinical and programmatic aspects, the article strives to enhance understanding of this significant pediatric illness. Strategies for strengthening prevention, timely case management and antibiotic stewardship based on identified research gaps are also proposed. Overall, the review intends to serve as a consolidated evidence base supporting enhanced multisectoral action against pneumonia, a leading cause of child mortality worldwide.

Methods

A systematic search strategy was employed to identify published literature on pneumonia in children below five years of age. Electronic databases including PubMed, MEDLINE, EMBASE, Cochrane Library, Web of Science and Global Health were searched using relevant keyword combinations focused on childhood pneumonia, epidemiology, etiology, clinical presentation, diagnosis, treatment guidelines, case management, risk factors, prevention strategies and developing countries. Primary literature published in English between 2000-2022 were considered to encompass recent insights. Hand searches of reference lists complemented the electronic database search.

Additional sources included pneumonia reports, operational guidelines and program reviews issued by leading public health agencies like WHO, UNICEF, Gates Foundation, CDC and country-level health ministries. Websites of non-profit organizations focused on pediatric pneumonia research and management such as PATH, PMNCH and Saving Lives at Birth were reviewed. Surveillance bulletins capturing epidemiological trends from high burden countries provided valuable epidemiological updates. Together these sources formed the basis for a comprehensive narrative synthesis.

Data on incidence, disease burden, risk factors, etiological agents, clinical presentations and case management practices were abstracted and analyzed. Regional and temporal insights on evolving priorities were charted. Evidence concerning diagnostic approaches, treatment strategies and prevention recommendations endorsed by agencies were compiled. Research gaps identified by subject matter experts and academia were incorporated to contextualize future directions. Due considerations were made regarding the variable quality and generalizability of available data. Summaries of key findings were drawn from both individual studies and consolidated reports, balanced against associated limitations and knowledge gaps.

Results

The results are presented under key domains including epidemiology, risk factors, etiology, clinical manifestations, diagnosis, treatment guidelines and prevention strategies.

IMPACT FACTOR (RESEARCH BIB) – 7,245
SPECIAL ISSUE, VOLUME 5, ISSUE 2, 2023

Epidemiology

Pneumonia accounts for approximately 15% of all under-five deaths globally with over 800,000 fatalities annually. Incidence is highest among neonates and children under 2 years. Developing regions bear over 95% of the global disease burden, with the highest rates in South Asia and sub-Saharan Africa. Pneumonia hospitalizations and outpatient visits constitute a substantial healthcare cost. Seasonal variations are seen, peaking in winter months in temperate regions.

Risk factors

Malnutrition is a major predisposing factor, impairing immunological defenses. Lack of exclusive breastfeeding in the initial six months deprives infants of critical antibodies. Indoor air pollution due to solid fuel use heightens exposure to respiratory pathogens. HIV infection, prematurity and low birth weight exacerbate disease susceptibility. Timely vaccination provides significant protection against the main bacterial etiologies.

Etiology

Streptococcus pneumoniae is the predominant causative organism overall, accounting for nearly 30% of cases globally. Respiratory syncytial virus (RSV) predominates in infants below 6 months. Haemophilus influenzae type b (Hib) was a major contributor prior to effective vaccines. Other viral and bacterial pathogens including influenza, parainfluenza, adenoviruses and Mycoplasma pneumoniae are associated geographically. Atypical and mixed etiologies are commonly seen.

Clinical manifestations

Typical features involve fever, cough, rapid or difficult breathing. Young infants may present with apnea, grunting respirations, nasal flaring or chest indrawing (signs of pneumonia severity). Identification of these danger signs prompts prompt medical management.

Diagnosis

Definitive diagnosis remains challenging given limited diagnostic capabilities especially in resource-poor settings. Key approaches include chest x-rays, viral antigen detection, bacterial culture and molecular diagnostics. However, given resource and feasibility constraints, clinical assessment forms the mainstay for diagnosis and treatment initiation in most settings. Point-of-care tests show promise but require scale up.

Treatment guidelines

The Integrated Management of Childhood Illness (IMCI) strategy endorsed by WHO emphasizes prompt initiation of oral antibiotics (amoxicillin) for non-severe pneumonia cases and injectable therapy (penicillin/ampicillin) for severe and very severe illness, guided by clinical symptom screening. Oxygen therapy, fluid management and supportive care are vital components.

Prevention strategies

Multifaceted approaches are needed including universal vaccination against Hib and pneumococcus, promotion of healthy home environments, exclusive breastfeeding, nutritional interventions, case

management of comorbidities like HIV/AIDS and health system preparedness for quality care provision. Expansion of integrated community case management also holds promise. Continued vaccine development targets additional viral etiologies and drug-resistant strains.

Discussion

This article aimed to consolidate prevailing evidence on multiple intersecting facets of pneumonia affecting children below school age globally. The review highlighted pneumonia as a leading threat to early childhood survival worldwide, despite progress enabled through vaccines, improved case management and expanded healthcare access in certain regions. Significant inter and intra-country disparities emphasize the need for context-specific analyses to address remaining prevention and control deficiencies.

Notably, malnutrition emerged as an overarching risk factor for pneumonia worsening across geographies, reinforced by suboptimal feeding practices like untimely supplementation. There was broad consistency regarding pneumococcus, RSV and bacterial Hib as top etiological pathogens in young children, warranting sustained vaccine prioritization. IMCI clinical algorithms endorsed by WHO were universally recognized as pragmatic diagnostic and treatment mainstays in resource-constrained settings due to feasibility and proven effectiveness for detecting non-severe pneumonia cases meriting oral antibiotics.

However, several knowledge gaps were also revealed. More precision is required concerning the attributable burden of non-bacterial viral triggers against which current interventions offer limited protection. Emerging antimicrobial resistance trends necessitate continual review of empiric treatment protocols. Low-cost point-of-care diagnostics deserve faster scale up to strengthen management and stewardship of antibiotic use. Operational barriers curb coverage and quality of community-level case management strategies warranting implementation research.

Conclusion

In conclusion, pneumonia continues to exact a heavy toll on the lives of young children, especially in developing nations. Concerted community engagement, optimized case management and enhanced prevention through vaccinations present opportunities to curb mortality. Addressing underlying determinants such as malnutrition through nutrition-sensitive social protection programs can reduce disease susceptibility. Widespread scale-up of proven cost-effective interventions tailored to local needs is imperative to accelerate progress against this important yet manageable cause of childhood morbidity and mortality.

References

1. Nair, H., Nokes, D. J., Gessner, B. D., Dherani, M., Madhi, S. A., Singleton, R. J., & Scott, J. A. (2010). Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. The Lancet, 375(9725), 1545-1555.
2. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J. E., & Black, R. E. (2015). Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. The Lancet, 385(9966), 430-440.

3. Rudan, I., Boschi-Pinto, C., Biloglav, Z., Mulholland, K., & Campbell, H. (2008). Epidemiology and etiology of childhood pneumonia. Bulletin of the World Health Organization, 86, 408-416.
4. Simões, E. A., & Sá-Leão, R. (2017). Epidemiology, etiology, pathogenesis and clinical manifestations of community-acquired pneumonia. Pediatric pulmonology, 52, S4-S13.
5. Rudan, I., El Arifeen, S., Black, R. E., & Campbell, H. (2007). Childhood pneumonia and diarrhoea: setting our priorities right. The Lancet infectious diseases, 7(1), 56-61.
6. Shah, S., Bachur, R. G., Kim, D., & Harper, M. B. (2018). Symptoms and Examination Findings in Children With Pneumonia Caused by Respiratory Syncytial Virus, Influenza Virus, and Bacteria. The Pediatric infectious disease journal, 37(9), 914.
7. Levine, O. S., O'Brien, K. L., Deloria-Knoll, M., Murdoch, D. R., Feudtner, C., Driscoll, A. J., ... & Kotloff, K. L. (2012). The pneumonia etiology research for child health project: a 21st century childhood pneumonia etiology study. Clinical Infectious Diseases, 54(suppl_2), S93-S101.

Links:

1. WHO guidelines on management of possible serious bacterial infection in young infants when referral is not possible: https://www.who.int/publications/i/item/9789241548045
2. UNICEF report on progress and challenges in pneumonia prevention and control: https://www.unicef.org/media/113966/file/UNICEF-Pneumonia-report-2022.pdf

WORDLY
KNOWLEDGE