IQRO JURNALI , IQRO JOURNAL, ИКРО ЖУРНАЛИ
ISSN:2181-4341| wordlyknowledge.uz/index.php/iqro
Volume 23 Issue 01 AVGUST 2026

UDC: 577.175.5:616.89-008.441

DISTINCTIVE FEATURES OF CANCER CELL METABOLISM: METABOLIC REPROGRAMMING AS A STRATEGY FOR SURVIVAL AND PROLIFERATION

Qo‘shshayeva Mahliyo Bobirjonovna
3rd-year Student, Faculty of Medicine
Tashkent State Medical University
Tashkent, Uzbekistan

Abstract

Cancer is traditionally characterized by uncontrolled cellular proliferation, but malignant transformation is accompanied by an equally profound reorganization of cellular metabolism. Cancer cells modify the utilization of glucose, amino acids, lipids, and mitochondrial substrates to satisfy the extraordinary demands of continuous growth. The classical Warburg effect, characterized by increased glucose consumption and lactate production despite the availability of oxygen, represents one of the most recognized metabolic features of malignant cells. However, contemporary understanding indicates that cancer metabolism is substantially more complex and adaptable than aerobic glycolysis alone. Tumor cells can combine glycolysis with mitochondrial oxidative phosphorylation, increase glutamine utilization, remodel lipid synthesis, alter amino acid metabolism, and strengthen antioxidant systems. These adaptations provide not only energy but also the molecular building blocks required for nucleic acid, protein, and membrane synthesis. Furthermore, metabolic products released by cancer cells can reshape the tumor microenvironment and influence immune-cell activity. This article examines the major biochemical characteristics of cancer cell metabolism and emphasizes metabolic plasticity as a central mechanism of tumor survival. Understanding these adaptations may contribute to the development of more selective metabolic interventions and personalized therapeutic strategies in oncology.

Keywords: cancer metabolism, metabolic reprogramming, Warburg effect, glycolysis, glutamine metabolism, mitochondrial metabolism, lipid metabolism, metabolic plasticity, tumor microenvironment, oxidative stress.

Introduction

Cancer development involves much more than the uncontrolled multiplication of cells. A malignant cell must continuously obtain energy, synthesize new cellular structures, maintain redox balance, tolerate environmental stress, and compete with surrounding cells for limited nutrients. These demands require profound changes in the way nutrients are acquired and metabolized.

Normal differentiated cells generally regulate their metabolic activity according to physiological requirements. Cancer cells, in contrast, frequently reorganize metabolic pathways in order to support persistent proliferation. This phenomenon is known as **metabolic reprogramming** and is now considered one of the fundamental characteristics of cancer biology.

The historical basis for this concept was established by Otto Warburg, who observed that tumor tissues frequently consumed glucose at high rates and produced large amounts of lactate even when oxygen was available. This phenomenon, subsequently termed the Warburg effect, challenged the traditional assumption that oxygen availability necessarily determines whether cells rely on glycolysis or mitochondrial respiration.

Modern research has demonstrated that the Warburg effect should not be interpreted as a complete replacement of mitochondrial metabolism. Many tumors retain highly active mitochondria and can rely considerably on oxidative phosphorylation. Instead, malignant cells demonstrate remarkable metabolic flexibility, switching between metabolic pathways according to their genetic characteristics and microenvironmental conditions.

Consequently, cancer metabolism is best understood as a dynamic network that simultaneously supports energy production, biosynthesis, redox regulation, cellular signaling, and adaptation to stress.

Purpose of the Study

The purpose of this study is to examine the distinctive metabolic characteristics of cancer cells and to explain how metabolic reprogramming contributes to malignant cell proliferation, survival, and adaptation to unfavorable environmental conditions. Particular attention is given to alterations in glucose metabolism, the Warburg effect, mitochondrial oxidative phosphorylation, glutamine and amino acid metabolism, lipid biosynthesis, redox homeostasis, and metabolic interactions within the tumor microenvironment.

The study also aims to emphasize the concept of metabolic plasticity as a fundamental feature of cancer biology. By integrating these metabolic pathways into a single framework, the study seeks to demonstrate how cancer cells redirect nutrients not only toward ATP production but also toward biosynthesis, antioxidant defense, signaling, and adaptation. A further objective is to consider how these metabolic characteristics may contribute to the identification of potential therapeutic vulnerabilities and the development of more selective metabolic approaches in cancer treatment.

Materials and Methods

This study was conducted as a narrative literature review. Relevant scientific articles and review papers on cancer cell metabolism were analyzed using academic databases and scientific publications. The literature search focused on cancer metabolic reprogramming, the Warburg effect, glycolysis, mitochondrial metabolism, glutamine and amino acid metabolism, lipid metabolism, redox homeostasis, and metabolic plasticity. Selected findings were critically reviewed and synthesized to provide an integrated overview of the distinctive metabolic characteristics of cancer cells.

Metabolic Reprogramming in Cancer Cells

Rapidly proliferating cancer cells have metabolic requirements that differ substantially from those of most differentiated cells. A dividing cell must duplicate its DNA, synthesize large quantities of proteins, construct new membranes, and increase its overall biomass. Therefore, ATP production alone cannot explain the metabolic behavior of tumors.

Cancer cells frequently redirect metabolic intermediates away from complete oxidation and toward anabolic pathways. Glucose-derived carbon, for example, can contribute to nucleotide, amino acid, and lipid synthesis. In this context, a metabolic pathway that appears inefficient from the perspective of ATP production may actually be advantageous because it provides intermediates required for cellular construction.

Metabolic reprogramming is influenced by oncogenic signaling pathways and tumor-suppressor dysfunction. Alterations involving pathways such as PI3K–AKT–mTOR, MYC, HIF signaling, and p53 can modify nutrient uptake and metabolic enzyme activity. Thus, genetic changes responsible for malignant transformation are closely connected with biochemical changes occurring inside the tumor cell.

This interaction creates a reciprocal relationship between genetics and metabolism: oncogenic signaling modifies metabolism, while metabolites themselves can influence signaling and epigenetic regulation.

The Warburg Effect and Enhanced Glycolysis

The Warburg effect remains one of the most recognizable metabolic characteristics of cancer cells. In this state, glucose is preferentially converted into pyruvate and subsequently lactate even when oxygen is sufficient for mitochondrial respiration.

At first glance, this strategy appears inefficient because oxidative phosphorylation can generate considerably more ATP from one glucose molecule. However, the biological objective of a proliferating tumor cell is not necessarily to maximize ATP yield per molecule of glucose. Instead, rapid glycolytic flux allows carbon intermediates to enter multiple biosynthetic pathways.

Increased glucose uptake is frequently supported by enhanced expression or activity of glucose transporters and glycolytic enzymes. Hexokinase, phosphofructokinase, pyruvate kinase, and lactate dehydrogenase are among the components that may contribute to increased glycolytic activity.

The conversion of pyruvate to lactate also regenerates NAD+, allowing glycolysis to continue rapidly. At the same time, lactate can accumulate in the tumor microenvironment and contribute to extracellular acidification.

Therefore, glycolysis in cancer should not be viewed simply as an alternative source of ATP. It functions as a **metabolic hub**, distributing glucose-derived carbon into pathways necessary for growth and adaptation.

Mitochondrial Metabolism in Malignant Cells

The classical interpretation of the Warburg effect sometimes led to the assumption that mitochondria become metabolically irrelevant in cancer. Contemporary evidence has demonstrated that this is not universally correct.

Cancer mitochondria remain involved in ATP generation, biosynthesis, redox regulation, apoptosis, and cellular signaling. Depending on tumor type and environmental conditions, some cancer cells may exhibit substantial dependence on oxidative phosphorylation.

The tricarboxylic acid cycle is particularly important because its intermediates can serve both energetic and biosynthetic functions. Citrate, α-ketoglutarate, oxaloacetate, and other intermediates can be redirected toward the synthesis of lipids, amino acids, and other cellular components.

Mitochondria can therefore be considered metabolic decision-making centers rather than merely energy-producing organelles. Their activity changes according to oxygen availability, nutrient supply, oncogenic signaling, and cellular stress.

This metabolic adaptability explains why inhibition of glycolysis does not necessarily eliminate all cancer cells. Some malignant populations can increase mitochondrial respiration or exploit alternative nutrient sources when glucose metabolism becomes restricted.

Glutamine Metabolism and Cellular Biosynthesis

Glutamine is another major nutrient supporting malignant growth. Although it is commonly described as a fuel, its biochemical significance extends considerably further.

Through glutaminolysis, glutamine can be converted into glutamate and subsequently into α-ketoglutarate, replenishing intermediates of the tricarboxylic acid cycle. This process can support mitochondrial metabolism and biosynthetic reactions.

Glutamine also provides nitrogen for the synthesis of nucleotides and other nitrogen-containing molecules. Because rapidly proliferating cells require large quantities of DNA and RNA precursors, nitrogen metabolism becomes particularly important during tumor expansion.

In addition, glutamine contributes indirectly to antioxidant defense by supporting pathways involved in glutathione synthesis. Consequently, glutamine can simultaneously support carbon metabolism, nitrogen metabolism, biosynthesis, and protection against oxidative stress.

Some tumors display a strong dependence on glutamine, sometimes described as glutamine addiction. However, this characteristic is not universal. Metabolic plasticity allows certain tumors to utilize alternative substrates when glutamine availability is reduced.

Amino Acid Metabolism and One-Carbon Metabolism

Amino acids serve several functions in cancer cells beyond protein synthesis. They can provide energy, carbon skeletons, nitrogen, and precursors for signaling molecules.

Serine and glycine metabolism is particularly important because these amino acids participate in one-carbon metabolism. This network contributes to nucleotide synthesis and methyl-group transfer reactions, linking amino acid metabolism with DNA synthesis and epigenetic regulation.

Alterations in amino acid metabolism may therefore influence both the physical growth of tumor cells and the regulation of gene expression.

Cancer cells can also modify amino acid transport systems to increase the intracellular availability of nutrients required for proliferation. Such changes demonstrate that metabolic reprogramming begins not only with intracellular enzymes but also with the mechanisms controlling nutrient entry into the cell.

Lipid Metabolism and Membrane Formation

Cell proliferation requires the production of new cellular membranes. For this reason, lipid metabolism represents another important component of tumor biology.

Cancer cells may increase de novo fatty acid synthesis even when extracellular lipids are available. Fatty acids are required for membrane formation, energy storage, signaling, and the production of lipid-derived regulatory molecules.

Enhanced lipid synthesis can therefore support the structural expansion of proliferating cells. At the same time, altered lipid metabolism can influence membrane composition and signaling pathways associated with malignant behavior.

This relationship between metabolism and cellular architecture illustrates an important principle: cancer cells do not simply require more energy; they require **more material from which to build new cells**.

Redox Homeostasis and Oxidative Stress

Rapid metabolism creates an additional biochemical challenge. Increased mitochondrial activity and other metabolic processes can elevate the production of reactive oxygen species.

Excessive reactive oxygen species can damage DNA, proteins, and lipids. Cancer cells therefore require mechanisms capable of maintaining redox homeostasis.

Antioxidant systems involving glutathione, NADPH, thioredoxin, and related pathways can help malignant cells tolerate elevated oxidative stress. This creates a metabolic paradox: tumor cells may generate increased levels of reactive oxygen species while simultaneously developing stronger antioxidant defenses.

The dependence of cancer cells on these protective systems creates a potential therapeutic opportunity. If antioxidant capacity is sufficiently disrupted, oxidative stress may exceed the cellular tolerance threshold and contribute to tumor-cell death.

Lactate and the Tumor Microenvironment

Lactate was traditionally regarded as an unwanted end product of glycolysis. In cancer biology, however, this interpretation is increasingly inadequate.

High glycolytic activity can result in substantial lactate production and export. Accumulation of lactate contributes to an acidic tumor microenvironment and can influence interactions between cancer cells and surrounding stromal and immune cells.

Importantly, lactate can also function as a metabolic substrate. Under certain conditions, tumor cells may take up extracellular lactate and utilize it for oxidative metabolism.

Thus, lactate can simultaneously represent a metabolic product, a signaling molecule, and a potential nutrient.

This observation illustrates the complexity of tumor metabolism: metabolites released by one population of cells may subsequently become resources for another population.

Metabolic Plasticity and Tumor Heterogeneity

One of the most important characteristics of cancer metabolism is its ability to change.

Tumors are not homogeneous structures composed of metabolically identical cells. Variations in oxygen concentration, nutrient availability, vascularization, genetic alterations, and interactions with stromal cells can create distinct metabolic states within the same tumor.

One group of cells may depend primarily on glycolysis, whereas another may rely more heavily on oxidative phosphorylation. Some cells may utilize glutamine extensively, while others can adapt to alternative carbon sources.

This phenomenon is known as **metabolic plasticity**.

From a therapeutic perspective, metabolic plasticity represents a major challenge. Blocking a single pathway may create selective pressure that encourages tumor cells to activate alternative pathways.

However, this same flexibility may also reveal combinations of metabolic dependencies that cannot be simultaneously bypassed. Identifying these metabolic “bottlenecks” could therefore become an important strategy in future cancer treatment.

Interaction Between Cancer Metabolism and the Immune System

Cancer metabolism does not occur independently of immunity. Tumor cells and immune cells compete for nutrients within the tumor microenvironment.

A rapidly growing tumor can consume substantial quantities of glucose and amino acids, potentially limiting their availability to immune cells. At the same time, metabolic products such as lactate can alter the local biochemical environment.

These changes may influence the activity of T cells, macrophages, and other immune populations. Consequently, tumor metabolism can contribute indirectly to immune evasion.

This connection between metabolism and immunity has created growing interest in combining metabolic interventions with immunotherapeutic approaches.

Rather than considering metabolism and immunity as separate therapeutic fields, modern cancer biology increasingly recognizes them as interconnected components of the same tumor ecosystem.

Therapeutic Perspectives

The distinctive metabolism of cancer cells provides several potential targets for therapy. Strategies under investigation include inhibition of glycolytic enzymes, glucose transport, glutamine utilization, lipid synthesis, mitochondrial respiration, amino acid transport, and antioxidant pathways.

Nevertheless, metabolic therapy presents a fundamental difficulty: normal cells also require these pathways. A successful treatment must therefore exploit differences in metabolic dependence between malignant and healthy tissues.

Another obstacle is metabolic plasticity. A tumor cell that loses access to one nutrient or pathway may compensate by increasing another. Consequently, combination therapy may be required to prevent metabolic escape.

Future approaches may increasingly focus on identifying the metabolic phenotype of an individual tumor before selecting treatment. Such precision-based strategies could improve therapeutic selectivity while minimizing damage to normal tissues.

Conclusion

Cancer cell metabolism represents a highly adaptable biochemical system designed to support continuous proliferation and survival under unfavorable conditions. The Warburg effect remains an important feature of this system, but it is only one part of a much broader metabolic network.

Cancer cells can increase glycolysis, retain active mitochondrial metabolism, exploit glutamine and other amino acids, remodel lipid synthesis, strengthen antioxidant defenses, and manipulate the metabolic environment surrounding the tumor. These pathways collectively provide energy, biosynthetic precursors, reducing equivalents, and signaling molecules.

The central characteristic of malignant metabolism is therefore not metabolic inefficiency but **metabolic flexibility**. Cancer cells continuously adjust their biochemical strategy according to nutrient availability, oxygen concentration, genetic alterations, and interactions with neighboring cells.

Understanding this flexibility may change the way metabolic therapies are designed. Instead of searching for one universal metabolic defect, future research may focus on identifying the

specific combinations of metabolic pathways upon which individual tumors are critically dependent.

Cancer metabolism can consequently be viewed as both a survival mechanism and a potential Achilles' heel. The same biochemical adaptations that allow malignant cells to survive may ultimately provide opportunities for more precise and effective therapeutic intervention.

References

1. Hammond, N. G., Cameron, R. B., & Faubert, B. (2024). Beyond glucose and Warburg: Finding the sweet spot in cancer metabolism models. *npj Metabolic Health and Disease, 2*, 11.
2. Jin, J., Byun, J. K., Choi, Y. K., et al. (2023). Targeting glutamine metabolism as a therapeutic strategy for cancer. *Experimental & Molecular Medicine, 55*, 706–715.
3. Ni, X., Lu, C. P., Xu, G. Q., et al. (2024). Transcriptional regulation and post-translational modifications in the glycolytic pathway for targeted cancer therapy. *Acta Pharmacologica Sinica, 45*, 1533–1555.
4. Chen, J., Cui, L., Lu, S., et al. (2024). Amino acid metabolism in tumor biology and therapy. *Cell Death & Disease, 15*, 42.
5. Demicco, M., Liu, X. Z., Leithner, K., et al. (2024). Metabolic heterogeneity in cancer. *Nature Metabolism*.
6. Tate, E. W., Soday, L., de la Lastra, A. L., et al. (2024). Protein lipidation in cancer: Mechanisms, dysregulation and emerging drug targets. *Nature Reviews Cancer, 24*, 240–260.

