IQRO JURNALI , IQRO JOURNAL, ИКРО ЖУРНАЛИ
ISSN:2181-4341| wordlyknowledge.uz/index.php/iqro
Volume 23 Issue 01 AVGUST 2026

UDC: 577.175.5:616.89-008.441

BIOCHEMICAL MECHANISMS OF HORMONAL REGULATION

Qo‘shshayeva Mahliyo Bobirjonovna
3rd-year Student, Faculty of Medicine
Tashkent State Medical University
Tashkent, Uzbekistan

Abstract

Hormonal regulation is a fundamental biochemical mechanism that coordinates the activity of cells, tissues, and organs and maintains physiological homeostasis. Hormones act as chemical messengers that transmit regulatory information from endocrine tissues to specific target cells. Their biological effects depend on the chemical properties of the hormone, the location of its receptor, intracellular signaling pathways, and the mechanisms responsible for terminating the hormonal response. Peptide and protein hormones generally interact with receptors located on the plasma membrane and activate intracellular signaling systems such as cyclic adenosine monophosphate, phosphoinositide signaling, calcium-dependent pathways, and protein kinase cascades. In contrast, steroid and thyroid hormones are predominantly lipid-soluble and regulate cellular functions through intracellular or nuclear receptors that influence gene transcription. Hormonal regulation is also characterized by feedback mechanisms, particularly negative feedback, which prevents excessive physiological responses and maintains endocrine stability. Disturbances in these biochemical processes may contribute to metabolic, endocrine, reproductive, and cardiovascular disorders. This article reviews the major biochemical mechanisms of hormonal action, receptor-mediated signaling, intracellular second messengers, regulation of gene expression, and feedback control, emphasizing their importance in maintaining physiological homeostasis.

Keywords: hormones, hormonal regulation, endocrine system, receptors, signal transduction, second messengers, gene expression, feedback regulation, homeostasis.

Introduction

The human organism requires continuous coordination between biochemical processes occurring in different cells and organs. Hormonal regulation provides one of the major mechanisms through which this coordination is achieved. Endocrine cells synthesize and release hormones that enter the circulation and subsequently interact with specific target tissues.

Hormones regulate a wide range of physiological processes, including metabolism, growth, development, reproduction, stress responses, electrolyte balance, and energy homeostasis. Their effects are highly specific because only cells containing appropriate receptors can respond to a particular hormonal signal.

The biochemical response to a hormone does not depend solely on the hormone itself. It involves a sequence of molecular events beginning with hormone recognition by a receptor and continuing through intracellular signaling pathways that ultimately modify cellular activity. Depending on the hormone and receptor type, these effects may occur within seconds through

modification of existing proteins or over a longer period through changes in gene transcription and protein synthesis.

Understanding the biochemical basis of hormonal regulation is therefore essential for explaining both normal physiology and the development of endocrine diseases.

Purpose of the Study

The purpose of this article is to describe the major biochemical mechanisms involved in hormonal regulation, including hormone-receptor interactions, intracellular signal transduction, second messenger systems, regulation of gene expression, and feedback mechanisms responsible for maintaining physiological homeostasis.

Materials and Methods

This article presents a narrative review of scientific literature concerning endocrine physiology, hormone biochemistry, receptor-mediated signaling, and intracellular regulatory mechanisms. Information from biomedical textbooks, scientific reviews, and research publications was analyzed and synthesized to describe the principal molecular mechanisms of hormonal action and their physiological significance.

Classification of Hormones

Hormones can be classified according to their chemical structure and physicochemical properties. The major groups include peptide and protein hormones, steroid hormones, amino acid derivatives, and several lipid-derived signaling molecules.

Peptide and protein hormones include insulin, glucagon, growth hormone, adrenocorticotropic hormone, and many hypothalamic and pituitary hormones. Because these molecules are generally water-soluble, they cannot readily cross the lipid bilayer of the plasma membrane. Their effects are therefore commonly mediated by receptors located on the cell surface.

Steroid hormones are synthesized primarily from cholesterol and include cortisol, aldosterone, estrogen, progesterone, and testosterone. Their lipid-soluble nature enables them to cross cellular membranes and interact with intracellular receptors.

Thyroid hormones are derived from tyrosine residues within thyroglobulin and possess lipophilic characteristics that allow them to act mainly through nuclear receptors. Catecholamines such as adrenaline and noradrenaline are also derived from amino acids but primarily exert their effects through membrane receptors.

The chemical structure of a hormone therefore strongly influences the mechanism through which it communicates with its target cell.

Hormone-Receptor Interaction

The first major step in hormonal signaling is the recognition of the hormone by its receptor. Receptors are specialized proteins capable of binding particular signaling molecules with high selectivity. Hormone binding induces conformational changes in the receptor and initiates a chain of intracellular events.

Membrane receptors are particularly important for peptide hormones and catecholamines. These receptors can be divided into several functional groups, including G-protein-coupled receptors, receptor tyrosine kinases, and receptors associated with intracellular kinases.

Intracellular receptors are characteristic of steroid and thyroid hormones. After entering the target cell, these hormones bind to receptors located in the cytoplasm or nucleus. The resulting

hormone-receptor complex can interact with specific DNA sequences and regulate transcription of selected genes.

Therefore, receptors function as molecular interfaces between extracellular hormonal signals and intracellular biochemical responses.

Second Messenger Systems

Many hormones produce their effects through intracellular second messengers. These molecules amplify the initial signal and allow a relatively small amount of hormone to produce a substantial cellular response.

One of the best-characterized pathways involves cyclic adenosine monophosphate (cAMP). When a hormone binds to a G-protein-coupled receptor, the receptor may activate a G protein that stimulates adenylate cyclase. Adenylate cyclase converts ATP into cAMP. The resulting increase in cAMP activates protein kinase A, which phosphorylates specific proteins and alters their activity.

Another important pathway involves phospholipase C. Activation of this system leads to the formation of inositol 1,4,5-trisphosphate and diacylglycerol. Inositol trisphosphate promotes the release of calcium ions from intracellular stores, while diacylglycerol contributes to protein kinase C activation. Together, these signaling molecules regulate numerous cellular processes.

Calcium itself can function as an important intracellular messenger. Changes in cytosolic calcium concentration can regulate enzymes, muscle contraction, secretion, and other cellular activities.

Receptor Tyrosine Kinase Signaling

Certain hormones and growth factors act through receptors possessing intrinsic or associated tyrosine kinase activity. Insulin is a major physiological example.

When insulin binds to its receptor, receptor activation results in phosphorylation of intracellular proteins and recruitment of signaling molecules. Important downstream pathways include the phosphatidylinositol 3-kinase/protein kinase B pathway and the mitogen-activated protein kinase pathway.

The first pathway is particularly important for metabolic effects of insulin. It contributes to glucose uptake, glycogen synthesis, lipid metabolism, and other processes involved in energy storage and utilization.

The complexity of these signaling networks allows one hormone to influence several cellular functions simultaneously while maintaining a coordinated physiological response.

JAK-STAT Signaling

Some peptide hormones and cytokines signal through the Janus kinase-signal transducer and activator of transcription pathway. In this mechanism, hormone binding activates receptor-associated Janus kinases, which phosphorylate specific STAT proteins.

Activated STAT proteins subsequently move into the nucleus, where they interact with regulatory DNA regions and influence gene transcription.

This mechanism demonstrates how a signal originating at the cell membrane can rapidly produce changes in gene expression without requiring a large number of intermediate signaling molecules.

Steroid Hormone Regulation of Gene Expression

Steroid hormones primarily influence cells by modifying gene transcription. Because they are lipid-soluble, they can pass through the plasma membrane and interact with intracellular receptors.

Following hormone binding, the receptor undergoes a structural change that enables interaction with specific regulatory regions of DNA. This interaction can increase or decrease transcription of selected genes.

The resulting changes in messenger RNA synthesis and protein production explain why steroid hormone effects are generally slower to develop than responses mediated by membrane receptors. However, transcriptional regulation can produce prolonged physiological effects.

Steroid hormones therefore provide an important connection between endocrine signaling and long-term modification of cellular phenotype.

Hormonal Regulation of Metabolism

Hormonal signaling plays a central role in maintaining energy balance. Insulin promotes glucose uptake and storage after food intake, while glucagon supports the mobilization of stored energy during periods of fasting.

Insulin stimulates glycogen synthesis, promotes anabolic processes, and suppresses excessive glucose production by the liver. Glucagon has largely opposing metabolic effects and stimulates glycogen breakdown and hepatic glucose production.

Adrenaline can rapidly increase the availability of energy substrates during acute stress. Cortisol contributes to longer-term metabolic adaptation by influencing carbohydrate, protein, and lipid metabolism.

The coordinated activity of these hormones enables the organism to adapt its metabolism to changes in nutritional status and physiological demands.

Feedback Regulation

Feedback mechanisms are essential for maintaining endocrine stability. Negative feedback occurs when the physiological effect produced by a hormone reduces further hormone secretion.

The hypothalamic-pituitary-adrenal axis provides an important example. Hypothalamic corticotropin-releasing hormone stimulates the pituitary secretion of adrenocorticotropic hormone, which promotes cortisol production by the adrenal cortex. Increased cortisol levels subsequently suppress the activity of the hypothalamus and pituitary, limiting further stimulation.

Positive feedback is less common but occurs in specific physiological situations. A classic example is the increase in oxytocin-mediated uterine contractions during labor, where the physiological response enhances further hormone release.

Feedback mechanisms prevent excessive hormonal activity and allow endocrine systems to respond dynamically to changing physiological conditions.

Termination of Hormonal Signals

Hormonal signaling must eventually be terminated to prevent continuous stimulation of target cells. Hormones may be metabolized in the liver, kidneys, or target tissues and subsequently eliminated from the body.

At the cellular level, signaling pathways can be attenuated through receptor desensitization, degradation of second messengers, dephosphorylation of activated proteins, or internalization of receptors.

For example, enzymes known as phosphodiesterases can break down cAMP, thereby reducing activation of protein kinase A. Protein phosphatases can remove phosphate groups from proteins activated by kinases.

These mechanisms ensure that hormonal responses remain proportional to physiological requirements.

Clinical Significance

Disruption of hormonal signaling can lead to significant pathological consequences. Abnormal hormone production, defective receptors, impaired intracellular signaling, or excessive hormonal stimulation may interfere with normal physiological functions.

Insulin resistance is an important example in which target tissues become less responsive to insulin signaling. Impaired insulin action contributes to abnormal glucose metabolism and plays a central role in the development of type 2 diabetes mellitus.

Disorders involving thyroid hormones, adrenal hormones, reproductive hormones, and growth-regulating hormones can similarly result from abnormalities at different levels of endocrine regulation.

Because hormonal pathways involve multiple interconnected molecular components, understanding their biochemical mechanisms is important for identifying potential therapeutic targets.

Discussion

Hormonal regulation represents a highly organized communication system in which extracellular signals are converted into specific biochemical responses. The diversity of receptor types and intracellular pathways allows hormones to produce effects ranging from rapid enzyme activation to long-term changes in gene expression.

The same hormone may produce different effects in different tissues because target cells can express distinct receptor subtypes and intracellular signaling proteins. In addition, the intensity of a hormonal response depends on receptor abundance, hormone concentration, intracellular messenger activity, and the mechanisms responsible for signal termination.

This complexity provides flexibility but also creates multiple points at which pathological changes may occur. Consequently, endocrine disorders can arise from abnormalities in hormone synthesis, secretion, transport, receptor function, signal transduction, or feedback regulation.

Conclusion

Hormonal regulation is an essential biochemical mechanism responsible for coordinating metabolism, growth, reproduction, stress adaptation, and maintenance of internal physiological stability. Hormones communicate with target cells through specific receptors and activate intracellular signaling pathways involving second messengers, protein kinases, transcription factors, and changes in gene expression.

Water-soluble hormones commonly produce rapid responses through membrane receptors, whereas lipid-soluble hormones frequently regulate cellular activity through intracellular receptors and transcriptional mechanisms. Feedback systems and signal termination pathways further ensure that endocrine responses remain controlled and appropriate.

A detailed understanding of these biochemical mechanisms is fundamental for explaining normal physiology and for understanding the molecular basis of endocrine and metabolic diseases.

References

1. Hall JE. *Guyton and Hall Textbook of Medical Physiology*. 14th ed. Elsevier; 2021.
2. Nelson DL, Cox MM. *Lehninger Principles of Biochemistry*. 8th ed. W.H. Freeman; 2021.
3. Berg JM, Tymoczko JL, Gatto GJ, Stryer L. *Biochemistry*. 9th ed. W.H. Freeman; 2019.
4. Melmed S, Auchus RJ, Goldfine AB, Koenig RJ, Rosen CJ. *Williams Textbook of Endocrinology*. 14th ed. Elsevier; 2020.
5. Alberts B, Johnson A, Lewis J, et al. *Molecular Biology of the Cell*. 7th ed. W.W. Norton; 2022.
6. Lodish H, Berk A, Kaiser CA, et al. *Molecular Cell Biology*. 9th ed. W.H. Freeman; 2021.
7. Brunton LL, Hilal-Dandan R, Knollmann BC. *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. 14th ed. McGraw Hill; 2022.
8. Boron WF, Boulpaep EL. *Medical Physiology*. 3rd ed. Elsevier; 2017.